arXiv:1212.0650v1 [cond-mat.mes-hall] 4 Dec 2012

Low-Frequency Admittance as Probe of Majorana Fermions

Christophe Mora[1] and Karyn Le Hur[2]
[1] *Laboratoire Pierre Aigrain, École Normale Supérieure,*
Université Paris 7 Diderot, CNRS; 24 rue Lhomond, 75005 Paris, France and
[2] *Center for Theoretical Physics, Ecole Polytechnique, CNRS, 91128 Palaiseau Cedex, France*

(Dated: December 5, 2012)

We investigate the admittance of a metallic quantum RC circuit with a spinful single-channel lead or equally with two conducting spin-polarized channels, as an ideal system to probe the dynamics of Majorana fermions. We address how the two-channel Kondo physics and its emergent Majoranas arise and we determine the universal crossover function describing the Fermi-liquid to non-Fermi liquid region. Remarkably, the same universal form emerges both at weak transmission and large transmission. We find that the charge relaxation resistance strongly increases in the non-Fermi liquid realm. Our findings can be measured using current technology assuming a large cavity.

PACS numbers: 73.63.Kv,72.15.Qm,71.10.Ay

The need for fast manipulation and readout of quantum coherent circuits, notably in the perspective of quantum computation, has been a strong motivation to investigate the dynamical response of nanoconductors [1]. Excited at frequencies ω in the quantum regime, $\hbar\omega \gg k_B T$, the systems evolve due to the intriguing interplay of correlation and quantum coherence effects. The quantum RC circuit [2], a quantum dot attached to a single lead and polarized by an external AC gate voltage, has emerged as the archetypical system for studying the dynamics of coherent circuits [3–11]. Recent experiments [12] on quantum hydrid structures combining microwave resonators [13] with semiconductor or nanotube quantum dots offer an alternative perspective to measure the admittance of quantum circuits. The existence of a quantized [14] charge relaxation (AC) resistance $R_q = h/(2e^2)$ in the quantum RC circuit has been shown [15] (and measured [2]) to originate from the Fermi liquid (FL) nature of low energy excitations where the elementary quasiparticles are non-interacting fermions. A deep connection [3] has also been drawn between the quantized resistance $R_q = h/e^2$ for large dots, the Shiba relation and the one-channel Kondo model in RC circuits [16] close to the charge degeneracy points.

In this paper, we investigate the non-Fermi liquid (NFL) situation where the elementary quasiparticles are Majorana fermions [17]. This description naturally applies to the quantum RC circuit with spinful (spin unpolarized) electrons and a large cavity (dot) [18–20] and is associated to the two-channel Kondo model [21]. It could be extended to the case of the helical edges of quantum spin Hall states [22–24]. The proof of the existence of Majorana fermions has engendered a spurt of experimental efforts in condensed-matter systems [25–27].

In the quantum RC circuit with two conducting (spin) modes, the singular points where subsequent charge states are degenerate produce NFL behavior at intermediate energies and for all gate voltages, while a crossover to FL physics is recovered at low energy [28]. The crossover energy scale vanishes at the charge degeneracy points. Here, we provide an analytical expression for charge fluctuations in the universal crossover bridging the FL and NFL regimes: the resistance now starts at $R_q = h/(2e^2)$ for a large cavity and rapidly increases with energy towards the NFL region. We believe that our full description in the NFL regime is also relevant for the charge dynamics of a topological superconductor hosting Majorana fermions at their edges [29].

The system under study comprises a large (metallic) quantum dot attached to a lead via a quantum point contact (QPC) with a single spin-unpolarized channel [18, 19, 30, 31]. The quantum RC circuit could be equally built at the helical edges of quantum spin-Hall insulators [11]. Electron confinement implies a charging energy $E_C = e^2/(2C_g)$, where C_g is the capacitance of the dot, and the interaction term $H_C = E_C(\hat{N} - N_0)^2$ in the Hamiltonian. N_0 is the dimensionless gate voltage and the operator $e\hat{N}$ gives the electron charge on the dot. This charge exhibits fluctuations, as a result of electrons tunneling in and out the lead, and defines the capacitance C_0 and the charge relaxation resistance R_q from the low frequency expansion [15]

$$e^2\chi_C(\omega) = C_0\left(1 + i\omega C_0 R_q\right) + \mathcal{O}(\omega^2) \tag{1}$$

of the retarded charge susceptibility $\chi_C(t-t') = i\theta(t-t')\langle[\hat{N}(t), \hat{N}(t')]\rangle$. Below, we address the extreme cases of weakly transmitting and almost transparent QPC. Although the mapping [16] to the two-channel Kondo model holds rigorously only at weak transmission and close to charge degeneracy, Matveev has argued that it influences the whole phase diagram. In fact, the effective model derived near perfect transmission can be shown [30] to be equivalent to the Abelian bosonization approach to the two-channel Kondo model at the Emery-Kivelson line [32], even in the presence of channel asymmetry [31].

We proceed and consider first an almost open dot with weak charge quantization. The model can be reduced to a one-dimensional form with coordinate x, the region $x < 0$

defining the lead and $x > 0$ the (infinite) dot. Electrons are weakly backscattered, with amplitude $r \ll 1$, at the boundary $x = 0$. In this regime, spin and charge excitations occur at well-separated energy scales, $r^2 E_C \ll E_C$, and the system is conveniently described using bosonization [33, 34] in the spin and charge sectors. The Hamiltonian takes the form $H = H_0 + H_{BS} + H_C$,

$$H_0 = \frac{v_F}{2\pi} \sum_{\alpha=c,s} \int_{-\infty}^{\infty} dx \left([\pi \Pi_\alpha(x)]^2 + [\partial_x \phi_\alpha(x)]^2 \right) \tag{2a}$$

$$H_{BS} = -\frac{2 v_F r}{\pi a} \cos\left(\sqrt{2}\phi_c(0)\right) \cos\left(\sqrt{2}\phi_s(0)\right) \tag{2b}$$

and the charge of the dot is given by $(e\sqrt{2}/\pi)\phi_c(0)$. The sequence of transformations on the spin fields that follows is inherited from Refs. [30, 35]: (i) an even/odd parity decomposition of the spin field decouples the odd part, since only $\phi_s(0)$ appears in $H_C + H_{BS}$. The even part is defined as $\phi_1(x) = (\phi_s(x) + \phi_s(-x))/\sqrt{2}$ for $x \geq 0$ (same for Π_s). (ii) one defines the chiral field $\phi(x) = \phi_1(|x|) + \pi \,\mathrm{sgn} x \int_0^{|x|} dy\, \Pi_1(y)$ unfolded onto the complete x-axis. (iii) In Eq. (2b), $\cos\left(\sqrt{2}\phi_s(0)\right) = \cos\phi(0)$ is refermionized with

$$\frac{1}{2\sqrt{\pi a}} e^{i\phi(x)} = \hat{a}\, \psi(x), \tag{3}$$

introducing explicitly the local Majorana fermion $\hat{a} = \hat{a}^\dagger$, $\hat{a}^2 = 1/2$, and a is a short-distance cutoff. In fact, only the imaginary part (chiral Majorana fermion) of the field $\eta(x) = (\psi(x) - \psi^\dagger(x))/(i\sqrt{2})$ appears in H_{BS} and the real part is decoupled.

Following these transformations, the action reads $S = S_F + S_c + S_{BS}$. S_F describes a free chiral Majorana fermion $\eta(x)$ and a free local Majorana fermion $\hat{a}$ with the unperturbed Green's functions $G_\eta^{(0)}(k,\omega) = (\omega - v_F k)^{-1}$ and $G_{\hat{a}}^{(0)}(\omega) = 1/\omega$,

$$S_c = \sum_{\omega_m} |\phi_c(\omega_m)|^2 \left(|\omega_m| + \frac{2E_C}{\pi} \right), \tag{4a}$$

$$S_{BS} = i r_0 \int_0^\beta d\tau\, \eta(\tau)\, \hat{a}(\tau)\, \cos\left(\sqrt{2}\phi_c(\tau) + \pi N_0\right), \tag{4b}$$

where $r_0 = 2 v_F\, r\, \sqrt{2/\pi a}$, $\eta(\tau) \equiv \eta(x=0,\tau)$ and $\phi_c(\tau) \equiv \phi_c(x=0,\tau)$. The charge part S_c has been obtained after integrating out the field $\phi_c(x)$ except at $x = 0$. The field ϕ_c has been shifted to transfer the gate voltage N_0 from the interaction term to the backscattering term.

As Eq. (4a) shows, the charge mode is massive and, below the charging energy E_c, $\phi_c(\tau)$ is pinned to zero. To leading order in r, a complete integration over the charge field ϕ_c results in $\phi_c = 0$ in S_{BS} and a renormalized r_0. In analogy with a barrier in a $g = 1/2$ Luttinger liquid [36], the backscattering term S_{BS} is a relevant perturbation that leads to NFL corrections. Fortunately, the refermionized action is an exactly solvable Majorana resonant level model [32]. The Lorentzian-shaped spectral function for the Majorana fermion $\hat{a}$ has a width $\Gamma = (8E_C\gamma/\pi^2)\, r^2 \cos^2(\pi N_0)$ such that $\Gamma \sim r^2 E_C \ll E_C$. Below the energy scale Γ, spin excitations are quenched and the NFL renormalization flow stops. The local Majorana fermion $\hat{a}$ is indeed screened and a crossover to a Fermi liquid regime is established [30].

In order to compute the charge susceptibility, we need to extend the analysis of Refs. [30, 37]. Instead of integrating the action Eq. (4a) over the whole charge mode ϕ_c, we stop the integration at the intermediate energy scale Λ such that $\Gamma \ll \Lambda \ll E_C$. By doing so, the massive part of the charge mode is almost fully integrated while we can still describe interactions between the spin and charge modes at energies $\sim \Gamma$. To leading order in $r \ll 1$, the factor $\cos\left(\sqrt{2}\phi_c(\tau) + \pi N_0\right)$ in Eq. (4a) is substituted by its average

$$\cos\left(\sqrt{2}\phi_c^l(\tau) + \pi N_0\right) e^{-\langle \phi_c(\tau)\phi_c(\tau)\rangle_{\varepsilon > \Lambda}} \tag{5}$$

where $\langle \phi_c(\tau)\phi_c(\tau)\rangle_{\varepsilon>\Lambda} \simeq (1/2)\ln(\omega_D \pi / 2E_C)$, $\omega_D = v_F/\gamma a$ is the high energy cutoff of the initial model required by linearization of the spectrum [38], and $\ln\gamma = \mathbf{C} \simeq 0.5772$ is the Euler constant. The field $\phi_c^l(\tau)$ contains the low energy part of the charge mode.

The controlled expansion of the cosine in Eq. (5) is then justified by the reduction of available energies, notably $\langle \phi_c^l(\tau)\phi_c^l(\tau)\rangle \sim (\Lambda/E_C)^2 \ll 1$. A second order expansion yields the effective action $S = S_F + S_c^l + S_{BS}^0 + S_{BS}^1 + S_{BS}^2$ where the Coulomb interaction part S_c^l keeps the structure of Eq. (4a) but with $|\omega_m| < \Lambda$. Furthermore,

$$\begin{aligned}
S_{BS}^0 &= i\sqrt{2\Gamma} \int_0^\beta d\tau\, \eta(\tau)\, \hat{a}(\tau), \\
S_{BS}^1 &= -2i\sqrt{\Gamma}\tan(\pi N_0) \int_0^\beta d\tau\, \phi_c^l(\tau)\, \eta(\tau)\, \hat{a}(\tau), \\
S_{BS}^2 &= -i\sqrt{2\Gamma} \int_0^\beta d\tau\, \left[\phi_c^l(\tau)\right]^2 \eta(\tau)\, \hat{a}(\tau).
\end{aligned} \tag{6}$$

In the rest of this paper, we shall consider the zero temperature limit $\beta \to \infty$ [39]. The action part $S_F + S_c^l + S_{BS}^0$ defines a non-interacting quadratic problem for which the unperturbed Green's functions are readily computed, for instance (retarded ones)

$$G_{\hat{a}}(\omega) = \frac{1}{\omega + i\Gamma}, \quad \langle \phi_c^l(\tau)\phi_c^l(0)\rangle_\omega = \frac{\pi^2/(4E_C)}{1 - i\omega\pi/(2E_C)}. \tag{7}$$

Here, $\langle ... \rangle_\omega$ refers to the Fourier transform. In the case of vanishing backscattering, $r = 0$, one obtains the susceptibility $\chi_C(\omega) = (2/\pi^2)\langle \phi_c^l(\tau)\phi_c^l(0)\rangle_\omega$ from Eq. (7). Compared with Eq. (1), this readily gives the quantum capacitance $C_0 = C_g$ and the charge relaxation resistance $R_q = h/(2e^2)$, half of the result of spinless electrons [3].

The correction $S_{BS}^1 + S_{BS}^2$ is included perturbatively, $\chi_C = K_0 + K_1 + K_2$, K_0 is the above result for $r = 0$, K_1 comes from the second order expansion of S_{BS}^1 and K_2 from the first order expansion of S_{BS}^2. After a lengthy but straightforward calculation, we obtain at zero temperature

$$e^2 K_0(\omega)/C_g = \alpha(\omega), \tag{8a}$$

$$e^2 K_1(\omega)/C_g = -8\gamma\, r^2 \sin^2(\pi N_0)\, \alpha(\omega)^2\, \Pi_{a\eta}(\omega), \tag{8b}$$

$$e^2 K_2(\omega)/C_g = -8\gamma\, r^2 \cos^2(\pi N_0)\, \alpha(\omega)^2\, \frac{\ln(E_C/\Gamma)}{2\pi} \tag{8c}$$

with $\alpha(\omega) = (1 - i\omega\pi/2E_C)^{-1}$. The logarithmic singularity is cutoff by the charging energy E_C. $\Pi_{a\eta}(\tau) = \langle \eta(\tau)\hat{a}(\tau)\eta\hat{a}\rangle$ is a polarization operator computed from the quadratic part of the action with the result

$$\begin{aligned} \Pi_{a\eta}(\omega) &= -(1/2\pi)\left[\ln(E_C/\Gamma) + \beta(\omega/\Gamma)\right], \\ \beta(x) &= -(1 + 2i/x)\ln(1 - ix). \end{aligned} \tag{9}$$

The function $\beta(\omega/\Gamma)$ describes the crossover between the FL and NFL responses for $\omega \ll \Gamma$ and $\omega \gg \Gamma$ respectively.

For $\omega \ll \Gamma$, we use the expansion

$$\beta(x) \simeq -2 + x^2/6 + ix^3/6 - 3x^4/20 + \ldots, \quad x \ll 1 \tag{10}$$

inserted in Eq. (8) and compare with Eq. (1) to extract C_0 and R_q. At vanishing frequency, the static susceptibility, and C_0, coincide precisely with Ref. [30]. Remarkably and similarly to the spinless case, we find no correction to the charge relaxation resistance $R_q = h/(2e^2)$ for $r \neq 0$ due to the absence of a linear term in Eq. (10). This result is confirmed by the Fermi liquid approach elaborated in Ref. [15] where lead electrons are coherently backscattered at $x = 0$ with a phase shift proportional to the static charge susceptibility [33]. One can then derive the Shiba relation and show that $R_q = h/(2e^2)$ for an arbitrary transmission of the QPC. Note that Γ vanishes at $N_0 = 1/2$ where the system is always a NFL.

We now turn to the opposite limit of weak transmission of the QPC. The system is adequately described [3, 16] by the tunnel Hamiltonian $H = H_0 + H_C + H_T$ where

$$H_T = t \sum_{k,k',s=\uparrow,\downarrow} \left(d_{ks}^\dagger c_{k's} + c_{k's}^\dagger d_{ks} \right) \tag{11}$$

transfers electrons between the (large) dot and the lead with operators c_k and d_k respectively; the index s refers, e.g., to the two spin polarizations. The free electron part reads $H_0 = \sum_{a=c,d,s} \varepsilon_k a_{ks}^\dagger a_{ks}$ for dot and lead electrons. H_T either decreases or increases the dot charge by one unit and thus does not commute with H_C. Far from charge degeneracy, the perturbative approach of Ref. [3] can be reproduced with an additional factor two that accounts for spin degeneracy. One readily obtains $R_q = h/(2e^2)$, again in agreement with the Fermi liquid picture. Perturbation theory however breaks down close to charge degeneracy $N_0 \simeq 1/2$ where NFL physics starts to play a role. In this region, the charge states other than 0 and 1 can be disregarded and a mapping to the two-channel Kondo model formulated [3, 16] where the two charge states are represented by a spin 1/2 with $\hat{N} = \frac{1}{2} + S_z$. The vicinity to charge degeneracy $h_0 = E_C(1 - 2N_0)$ defines a local magnetic field coupled to S_z. Our study of charge fluctuations is then translated to a study of the local spin susceptibility in the two-channel Kondo model.

For $h_0 \ll T_K$, where T_K is the Kondo temperature, two regimes have been identified [21] in the renormalization group (RG) analysis: NFL properties dominate for frequencies (energies) $h_0^2/T_K < \omega < T_K$ while a FL response is obtained at smaller frequencies $\omega < h_0^2/T_K$. The crossover is investigated analytically using the SO(8) representation [40] of the two-channel Kondo model, which provides a simple description of the NFL fixed point [41, 42]. The bulk fermions, with two spin species and two channels, have a non-local representation in terms of eight chiral Majorana fermions. With no impurity, the free Hamiltonian reads

$$H_0^{\mathrm{K}} = \frac{-iv_F}{2} \sum_{j=1}^{8} \int_{-\infty}^{+\infty} dx\, \chi_j(x) \partial_x \chi_j(x). \tag{12}$$

The Majorana fermions $\chi_{1,2,3}$ generate the spin current, $\chi_{4,5,6}$ the flavor current and χ_7, χ_8 the charge current.

In the presence of the Kondo impurity coupled only to the spin current, the NFL infrared fixed point is simply characterized by the twisted boundary conditions $\chi_j(0^-) = -\chi_j(0^+)$ for $j = 1, 2, 3$. Absorbing this $\pi/2$ phase shift into a redefinition of the fields, $\chi_{1,2,3}(x) \to \mathrm{sgn}(x)\chi_{1,2,3}(x)$, one recovers the free Hamiltonian form Eq. (12) also at the infrared fixed point. A finite local magnetic field h_0 destabilizes this fixed point with the relevant perturbation [41] $H_{\mathrm{b}} = i(h_0/\sqrt{T_K/v_F})\chi_1\hat{a}$ of scaling dimension 1/2, where $\chi_1 = \chi_1(0)$. The local Majorana fermion $\hat{a}$ describes the residual impurity spin. The Hamiltonian $H_{IR} = H_0^K + H_b$ is equivalent to the two-dimensional Ising model with a boundary magnetic field, a correspondence that has been used to calculate the one-body Green's function along the FL to NFL crossover [42]. For energies $\omega \ll h_0^2/T_K$, the relevant boundary term H_{b} restores the continuity of $\chi_1(x)$ at $x = 0$. We recover a Fermi liquid as the even number of twisted fields (χ_2 and χ_3) indicates.

Quite generally, the impurity spin can be expanded over the different operators allowed by conformal field theory (CFT). At low energy, the leading term is

$$S_z = i\sqrt{\frac{v_F}{T_K}}\, \chi_1 \hat{a}, \tag{13}$$

in accordance with H_b. The spin susceptibility $\chi_s(\tau) = -(v_F/T_K)\,\langle \chi_1(\tau)\hat{a}(\tau)\chi_1\hat{a}\rangle$ is obtained by noting the equivalence between the Hamiltonian H_{IR} and the quadratic action $S_F + S^0_{BS}$, derived in the large transparency case. We identify $\eta = \chi_1$ and $2\Gamma = h_0^2/T_K$ and find

$$\chi_{s0}(\omega) = \frac{1}{2\pi T_K}\left[\ln\left(\frac{T_K}{\Gamma}\right) + \beta(\omega/\Gamma)\right], \tag{14}$$

describing the FL-NFL crossover. At large frequency $\omega \gg \Gamma$, the spin susceptibility exhibits NFL features

$$\chi_{s0}(\omega) = \frac{1}{2\pi T_K}\left[\ln\left(\frac{T_K}{\omega}\right) + \frac{i\pi}{2}\right], \tag{15}$$

in agreement with the prediction of Conformal Field Theory [21] and Abelian bosonization [32]. The absence of a linear term in Eq. (10) requires, for the calculation of R_q, to include the leading irrelevant perturbation to H_{IR},

$$\delta H = \frac{2\pi v_F^{3/2}}{\sqrt{T_K}}\,\chi_1\chi_2\chi_3\hat{a}. \tag{16}$$

The only linear in frequency correction to the spin susceptibility comes from the vertex correction, shown Fig.1 a), $\delta\chi_s(\omega) = \chi_{s0}(\omega)\Pi_e(\omega)\chi_{s0}(\omega)$ where $\Pi_e(\omega)$ is the electron-hole pair susceptibility. At zero temperature, $\Pi_e(\omega) = i\pi\omega$, interpreted as the dissipation produced by electron-hole excitations. Expanding the spin susceptibility $\chi_s = \chi_{s0} + \delta\chi_s$ to linear order in ω, we arrive at the Shiba relation

$$\text{Im}\chi_s(\omega) = \hbar\pi\omega\,\chi_s(0)^2, \tag{17}$$

equivalent to the charge relaxation resistance $R_q = h/(2e^2)$. This result confirms the validity of the Fermi liquid picture [15] also at low transmission.

Writing $\chi_s = \chi_1 + i\chi_2$, a frequency-dependent charge relaxation resistance can be defined from $R_q(\omega) = (\hbar/e^2)\chi_2(\omega)/(\omega\chi_1^2(\omega))$. At frequencies $\omega \ll T_K$, we extract the universal form

$$\frac{R_q(\omega)}{h/e^2} = A\,\Phi\left(\frac{\omega}{\Gamma}\right) = A\,\frac{\Gamma}{\omega}\frac{\text{Im}\beta(\omega/\Gamma)}{B + \text{Re}\beta(\omega/\Gamma)}, \tag{18}$$

with $A = T_K/\Gamma$. The dimensionless function $\Phi(x)$ is plotted in Fig.1 b) for different values of $B = \ln(T_K/\Gamma)$.

Remarkably, the same scaling form involving the function Φ is obtained in the opposite limit of weak backscattering. In the scaling limit where $N_0 \to 1/2$ and $\omega \ll E_C$, one has $\alpha(\omega) \simeq 1$, $\sin(\pi N_0) \simeq 1$, $K_2 \simeq 0$ and $K_0 \ll K_1$ in Eqs. (8). As a result, one recovers Eq. (18) for the charge relaxation resistance R_q with $B = \ln(E_C/\Gamma)$ and $A = E_C/(4\gamma\, r^2\,\Gamma)$.

To summarize briefly, we have shown that the presence of Majoranas in a quantum RC circuit results in a subtle charge dynamics which can, in principle, be revealed using current technology [39]. The FL to NFL crossover produces a visible increase of the charge relaxation resistance which can be probed through admittance measurements. We anticipate that a similar crossover should also be relevant for a superconducting wire supporting Majorana fermions at his edges [25, 26, 29]. Majorana fermions can also be detected using an optical cavity [43]. Other interesting directions concern the role of an asymmetry between channels which can be engineered through Zeeman effects for example [44].



FIG. 1. (a) Vertex correction to the spin susceptibility χ_s to second order in δH. (b) Dimensionless function $\Phi(\omega/\Gamma)$ for $B = 5, 7, 10$ (solid, dotted and dashed lines). We note that the quantized result $R_q = h/2e^2$, recovered at zero frequency, is not visible in this plot obtained in the scaling limit $A \gg 1$.

We acknowledge discussions with I. Affleck, P. Dutt, M. Filippone, B. Horovitz, P. Le Doussal, Z. Ristivojevic, T. Schmidt and E. Sela. CM thanks L. Glazman for fruitful discussions and comments. KLH acknowledges support from DOE under the grant DE-FG02-08ER46541.

---

[1] R. J. Schoelkopf, P. Wahlgren, A. A. Kozhevnikov, P. Delsing, and D. E. Prober, Science **280**, 1238 (1998)

[2] J. Gabelli, G. Fève, J.-M. Berroir, B. Plaçais, A. Cavanna, B. Etienne, Y. Jin, and D. C. Glattli, Science **313**, 499 (2006)G. Fève, A. Mahé, J.-M. Berroir, T. Kontos, B. Plaçais, D. C. Glattli, A. Cavanna, B. Etienne, and Y. Jin, *ibid.* **316**, 1169 (2007)J. Gabelli, G. Fève, J. Berroir, and B. Plaçais, arXiv:1210.2874(2012)

[3] C. Mora and K. Le Hur, Nature Phys. **6**, 697 (2010)

[4] S. E. Nigg, R. López, and M. Büttiker, Phys. Rev. Lett. **97**, 206804 (2006)

[5] Y. I. Rodionov, I. S. Burmistrov, and A. S. Ioselevich, Phys. Rev. B **80**, 035332 (2009)

[6] Y. Hamamoto, T. Jonckheere, T. Kato, and T. Martin, Phys. Rev. B **81**, 153305 (2010)

[7] J. Splettstoesser, M. Governale, J. König, and

M. Büttiker, Phys. Rev. B **81**, 165318 (2010)

[8] Y. Etzioni, B. Horovitz, and P. Le Doussal, Phys. Rev. Lett. **106**, 166803 (2011)

[9] M. Lee, R. López, M.-S. Choi, T. Jonckheere, and T. Martin, Phys. Rev. B **83**, 201304 (2011)

[10] C. Grenier, R. Hervé, G. Fève, and P. Degiovanni, Mod. Phys. Lett. B **25**, 1053 (2011)

[11] I. Garate and K. Le Hur, Phys. Rev. B **85**, 195465 (2012)

[12] M. R. Delbecq, V. Schmitt, F. D. Parmentier, N. Roch, J. J. Viennot, G. Fève, B. Huard, C. Mora, A. Cottet, and T. Kontos, Phys. Rev. Lett. **107**, 256804 (2011)S. J. Chorley, J. Wabnig, Z. V. Penfold-Fitch, K. D. Petersson, J. Frake, C. G. Smith, and M. R. Buitelaar, *ibid.* **108**, 036802 (2012)T. Frey, P. J. Leek, M. Beck, A. Blais, T. Ihn, K. Ensslin, and A. Wallraff, *ibid.* **108**, 046807 (2012)T. Frey, P. J. Leek, M. Beck, J. Faist, A. Wallraff, K. Ensslin, T. Ihn, and M. Büttiker, Phys. Rev. B **86**, 115303 (2012)K. D. Petersson, L. W. McFaul, M. D. Schroer, M. Jung, J. M. Taylor, A. A. Houck, and J. R. Petta, Nature **490**, 380 (2012)H. Toida, T. Nakajima, and S. Komiyama, arXiv:1206.0674(2012)

[13] A. Wallraff, D. I. Schuster, A. Blais, L. Frunzio, R. S. Huang, J. Majer, S. Kumar, S. M. Girvin, and R. J. Schoelkopf, Nature **431**, 162 (2004)

[14] M. Büttiker, A. Prêtre, and H. Thomas, Phys. Rev. Lett. **70**, 4114 (1993)M. Büttiker, H. Thomas, and A. Prêtre, Phys. Lett. A **180**, 364 (1993)A. Prêtre, H. Thomas, and M. Büttiker, Phys. Rev. B **54**, 8130 (1996)

[15] M. Filippone, K. Le Hur, and C. Mora, Phys. Rev. Lett. **107**, 176601 (2011)M. Filippone and C. Mora, Phys. Rev. B **86**, 125311 (2012)

[16] L. Glazman and K. A. Matveev, Zh. Eksp. Teor. Fiz. **98**, 1834 (1990), [*Sov. Phys. JETP* **71,** 1031 (1990)]K. A. Matveev, *ibid.* **99**, 1598 (1990), [*Sov. Phys. JETP* **72,** 892 (1991)]

[17] E. Majorana, Nuovo Cimento **14**, 171 (1937)

[18] D. Berman, N. B. Zhitenev, R. C. Ashoori, and M. Shayegan, Phys. Rev. Lett. **82**, 161 (1999)

[19] K. W. Lehnert, B. A. Turek, K. Bladh, L. F. Spietz, D. Gunnarsson, P. Delsing, and R. J. Schoelkopf, Phys. Rev. Lett. **91**, 106801 (2003)

[20] E. Lebanon, A. Schiller, and F. B. Anders, Phys. Rev. B **68**, 041311 (2003)

[21] D. L. Cox and A. Zawadowski, Adv. Phys. **47**, 599 (1998)

[22] C. L. Kane and E. J. Mele, Phys. Rev. Lett. **95**, 226801 (2005)

[23] B. A. Bernevig and S.-C. Zhang, Phys. Rev. Lett. **96**, 106802 (2006)

[24] M. Koenig, S. Wiedmann, C. Bruene, A. Roth, H. Buhmann, L. W. Molenkamp, X.-L. Qi, and S.-C. Zhang, Science **336**, 1003 (2007)

[25] V. Mourik, K. Zuo, S. M. Frolov, S. R. Plissard, E. P. A. M. Bakkers, and L. P. Kouwenhoven, Science **336**, 1003 (2012)

[26] L. P. Rokhinson, X. Liu, and J. K. Furdyna, arXiv:1204.4212(2012)

[27] J. R. Williams, A. J. Bestwick, P. Gallagher, S. S. Hong, Y. Cui, A. S. Bleich, J. G. Analytis, I. R. Fisher, and D. Goldhaber-Gordon, Phys. Rev. Lett. **109**, 056803 (2012)

[28] I. Affleck, A. W. W. Ludwig, H.-B. Pang, and D. L. Cox, Phys. Rev. B **45**, 7918 (1992)

[29] A. Golub and E. Grosfeld, arXiv:1206.0958(2012)

[30] K. A. Matveev, Phys. Rev. B **51**, 1743 (1995)

[31] K. Le Hur and G. Seelig, Phys. Rev. B **65**, 165338 (2002)

[32] V. J. Emery and S. Kivelson, Phys. Rev. B **46**, 10812 (1992)D. G. Clarke, T. Giamarchi, and B. Shraiman, *ibid.* **48**, 7070 (1993)A. M. Sengupta and A. Georges, *ibid.* **49**, 10020 (1994)

[33] I. L. Aleiner and L. I. Glazman, Phys. Rev. B **57**, 9608 (1998)

[34] L. I. Glazman, F. W. J. Hekking, and A. I. Larkin, Phys. Rev. Lett. **83**, 1830 (1999)

[35] I. L. Aleiner and A. V. Andreev, Phys. Rev. Lett. **81**, 1286 (1998)

[36] C. L. Kane and M. P. A. Fisher, Phys. Rev. B **46**, 15233 (1992)

[37] A. Furusaki and K. A. Matveev, Phys. Rev. Lett. **75**, 709 (1995)Phys. Rev. B **52**, 16676 (1995)

[38] K. Flensberg, Phys. Rev. B **48**, 11156 (1993)

[39] S. Amasha, I. G. Rau, M. Grobis, R. M. Potok, H. Shtrikman, and D. Goldhaber-Gordon, Phys. Rev. Lett. **107**, 216804 (Nov 2011), `http://link.aps.org/doi/10.1103/PhysRevLett.107.216804`

[40] J. M. Maldacena and A. W. W. Ludwig, Nucl. Phys. B **506**, 565 (1997)J. Ye, Phys. Rev. Lett. **79**, 1385 (1997)

[41] E. Sela and I. Affleck, Phys. Rev. B **79**, 125110 (2009)J. Malecki, E. Sela, and I. Affleck, **82**, 205327 (2010)A. K. Mitchell, E. Sela, and D. E. Logan, Phys. Rev. Lett. **108**, 086405 (2012)

[42] E. Sela, A. K. Mitchell, and L. Fritz, Phys. Rev. Lett. **106**, 147202 (2011)A. K. Mitchell and E. Sela, Phys. Rev. B **85**, 235127 (2012)

[43] T. L. Schmidt, A. Nunnenkamp, and C. Bruder, arXiv:1211.2201(2012)

[44] P. Dutt, T. L. Schmidt, C. Mora, and K. Le Hur, to appear on ArXiv(2012)